

IFTAK AN INNOVATIVE TECHNIQUE IN FISTULA IN ANO-A CASE STUDY

¹Dr Amit Kumar ²Dr Anchal Bilyan

¹Assistant Professor, Shalya Tantra Dept. Uttaranchal Ayurvedic College, Dehradun (U.K)

²Assistant Professor, Kayachikitsa Dept. Uttaranchal Ayurvedic College, Dehradun (U.K)

ABSTRACT

Fistula in ano is an infective disease of anal canal usually caused by crypto glandular infection of anal crypts. In fistula in ano there is an abnormal communication between two epithelial surfaces and the track is lined by unhealthy granulation tissue. It can be correlated with *Bhagandara* described in Ayurveda texts. The present case study was conducted on a male patient aged 56 yrs. Who was suffering from Fistula in ano for last 3 yrs. IFTAK (Interception of fistulous tract with application of *Kshara sutra*) technique of treatment was adopted. Daily dressings were done with *jatyaditaila*. Patient cured completely within 4 weeks of treatment. There was no side effect or complication after regular followup of 2 months. The treatment undertaken was very effective as it reduced seton placement in smaller track and also reduced healing time of the wound. The treatment techniques proved to be very effective and convective treatment option.

KEYWORDS: Bhagandara, Fistula in ano, IFTAK, Jatyaditaila, Kshara sutra

INTRODUCTION

Fistula in ano is a notorious disease of anal canal. It is mainly caused by crypto glandular infection of anal crypt. A fistula-in-ano, or anal fistula, is a chronic abnormal communication, usually lined to some degree by unhealthy granulation tissue, which runs outwards from the ano-rectal lumen (the internal opening) to an external opening on the skin of the perineum or buttock (or rarely, in women, to the vagina). Anal fistulae may be found in association with specific conditions, such as Crohn's disease, tuberculosis, lymphogranuloma venereum, actinomycosis¹. In Ayurveda *Bhagandara* can be correlated with fistula in ano. *Bhagandara* is painful condition of anal and perianal region with discharge of pus². According to Vagbhata, *Bhagandara*

The disease which causes *darana* (deformity) in and around *bhaga* (pubic region, perineum, vaginal region, and genital area), *guda* (anal region) and *basti* (urinary bladder) is called *Bhagandara*. When the blister remains unripe (not suppurated) it is called *Pidaka*, when the same gets suppurated it will be called *Bhagandara*⁴. The present case study was done on a case of *Arshobhagandara*. An anal fistula (also commonly called fistula-in-ano) is frequently the result of a previous or current anal abscess. This occurs in up to 50% of patients with abscesses. Normal anatomy includes small glands just inside the anus. The fistula is the tunnel that forms under the skin and connects the clogged infected glands to an abscess. A fistula can

be present with or without an abscess and may connect just to the skin of the buttocks near the anal opening⁵.

Technique

IFTAK (Interception of fistulous tract with application of Ksharasutra)⁶

IFTAK is also known as BHU technique of treatment of fistula in ano. The technique was developed by Dr M. Sahu, (Professor, Department of Shalya Tantra, faculty of Ayurveda, Banaras Hindu University, Varanasi, UP. India) and is being practiced for treating complex and recurrent fistula in ano in Banaras Hindu University, Varanasi since 2007.

In this technique, proximal part of fistulous track is intercepted at the level of external sphincter along with the application of *Ksharasutra* from site of interception to the infected crypt in anal canal. This is aimed at to eradicate the infected anal crypt with no or minimal damage to anal sphincters by using *ksharasutra* (medicated seton). Use of *ksharasutra* causes extensive fibrosis and favors proper healing which reduces the chances of recurrence⁷.

CASE:

The present study was conducted on a male patient aged 56yrs who came to UAH (Uttaranchal Ayurvedic Hospital) with complaints of pain, swelling and pus discharge from anal canal for last 3 yrs. Patient was not a known case of Diabetes mellitus/Hypertension/Tuberculosis/Bronchial asthma. Patient was examined properly and diagnosed as Fistula in Ano with extero-internal haemorrhoids.

Short Clinical History

History of Present Illness

Patient was asymptomatic 3 yrs ago. Then he gradually developed pain in perianal region, which gradually increased in intensity. After 1-2 months, there was swelling in perianal area and subsequently discharge of pus from that region. Initially the pus discharge occurred during defecation, but after some time the discharge was at any time of the day. There was fever while having pus discharge. There was a lot of discomfort in sitting and daily routine work. Patient took some medications from nearby medical facility which lessen the local symptoms. After 4-5 months patient developed similar complaints. Patient had history of on and off occurrence of the previous symptoms. With these complaints patient was admitted in Uttaranchal Ayurvedic Hospital (UAH).

Family History

No history of similar complaint in the family

Personal History

- >Bowel- Constipation
- >Bladder- Regular
- >Sleep- Inadequate
- >Appetite- Normal
- >Addiction- No addiction

Physical Examination

Average built, no clubbing/cyanosis/icterus, no any lymphadenopathy observed.

Systemic Examination

- >Respiratory System
- Bilateral air entry- normal
- No any added sounds
- >Cardiovascular System
- S1, S2 – normal
- No murmur
- >Central Nervous system

Patient fully conscious, well oriented to time place & person

Local Examination

1. An external opening at 3 o'clock approx. 3.5cm away from anal verge with hyper granulated area.
2. Internal tender dimpling was noted at 6 o'clock below dentate line with tender perianal region at 6 o'clock. There was no dimpling at 3 o'clock.
3. On Syringing from external opening- Solution effused at 6 o'clock visualized with the help of slit proctoscope.
4. IInd degree extero internal haemorrhoids at 7 o'clock with a skin tag at 6 o'clock.

Investigations-

CBC- Within normal limits (WNL)

HIV I & II- Negative

RBS- 76 mg/dl

HBsAg- Non reactive

Treatment

Ksharasutra Treatment was planned with IFTAK (Interception of Fistulous track with application of *Ksharasutra*) Technique.

PROCEDURE- *Ksharasutra* therapy

>Procedure was done under local anaesthesia.(L.A)

>After painting and Draping Local anesthesia was given. Probing of fistulous track done which was communicating external opening at 3 o'clock and internal dimpling at 6 o'clock.

>External opening was opened and widened. A small incision was made at perianal area at 6 o'clock approx. 1.5 cm away from anal verge. Pus was drained from that opening. Approx. 4 ml pus drained.

>Metallic probe was introduced through the opening and taken out through internal opening. Primary threading was done.

>Haemorrhoidectomy of 7o'clock pile mass done after transfixation with *kshara sutra*.

>Antiseptic dressing done and packing done with *Jatyadi taila*.

>On next day *kshara* sutra was applied into the track replacing the primary threading.

>Patient was advised hot sitz bath daily dressing with *jatyadi taila*.

Oral medication- *Triphala guggulu* 2 tab. BD was given for 15 days.

Regular followup was advised with weekly *ksharasutra* change.

Follow UP

Regular follow up were done to assess the progress of the disease. The pus discharge was fluent in 1st 3 days. After that it reduced and completely disappeared after 7-10 days. The wound at 3 o'clock gradually healed. After 4 *ksharasutra* changes which was done at weekly interval, the track at 6 o'clock was laid open and packing of *jatyaditaila* impregnated gauze was continued. In next followup after 7 days both the wound healed completely.

OBSERVATION

After treatment patient was observed on regular basis. Progress of treatment was noted. The fistula track healed in 4weeks completely with minimum scarring. Patient returned to normal daily routine work with no side effect or adverse effect after one month. There was no recurrence of the symptoms after 6 months of follow up

DISCUSSION

The treatment adopted to treat the fistula in ano proved to be very effective. It reduced the healing time and avoided conventional threading into a long thread communicating external and internal opening. The technique focused on the crypt, which was involved in

the pathogenesis of fistula in ano. Otherwise in conventional technique of *ksharsutra* therapy, a long track would have formed which requires a long time to heal and discomfort to the patient.

The secondary track/wound healed subsequently with the healing effect of *jatyaditaila*. *Triphla guggulu* worked as anti-infective preventing secondary infection as well as a mild laxative. The whole time period of treatment was around one month. Patient was on his daily routine work 5th day onwards.

CONCLUSION

This innovative technique is quite helpful in treating fistula in ano having distant external opening. In the conventional technique of threading it would have involved larger track communicating internal and external opening. But in IFTAK technique threading was done in much smaller track. It reduced healing time and postoperative scar and caused less irritation to the patient.

REFERENCES

1. Bailey & Love's Short Practice of Surgery, edited by Norman S. Williams, Christopher J.K. Bulstrode, P. Ro-nan O'Connell, Hodder Arnold publication, 25th Edition, Anus and Anal Canal, *Fistula-in-ano*, 1262.
2. Shashtri A.D., Sushrut Samhita in Ayurveda Tatva Sandipika, chaukhamba Sanskrit sansthan, nidansthana chapter Su. Ni. 4/4, p-244
3. Tripathi Bhramanand Astanga hridayam, nirmalahindi commentary, chaukhamba Sanskrit sansthan Delhi, uttarsthan chapter- 28/5, p-1092
4. Shashtri A.D., Sushrut Samhita in Ayurveda Tatva Sandipika, chaukhamba Sanskrit sansthan, nidansthanam chapter Su. Ni. 4/12, p-246

5. <https://www.fascrs.org/patients/disease-condition/abscess-and-fistula-expanded-information> dated 25.3.18

6. Rojanasakul A. LIFT procedure: a simplified technique for fistula-in-ano. Tech Coloproctol, 2009 Sep; 13(3): 237-40. Epub 2009 Jul 28.

7. M. Sahu, A manual on fistula in ano and kshara sutra therapy, published by NRC, Deptt. Of shalyatantra, IMS, BHU, first edition, 2015.

CORRESPONDING AUTHOR

Dr. Amit Kumar,

Assistant Professor, Dept. of Shalya Tantra
Uttaranchal Ayurvedic College, Dehradun
(U.K)

E-mail: amitnegiayurveda@gmail.com

Source of support: Nil,

Conflict of interest: None Declared

Cite this article as

Amit Kumar: IFTAK An Innovative
Technique In Fistula In Ano-A Case Study
ayurpub;III(2): 771-775



Fig I. Before Procedure



Fig II. After Procedure



Fig III. After 1 week



Fig IV. After 4 weeks